

VACCINE FOR PREVENTING PYTHIOSIS
IN HUMANS AND ANIMALS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. application Serial No. 09/082,232, filed May 20, 1998, now U.S. Patent No. 6,287,573, which is a division of Serial No. 08/895,940, filed July 17, 1997, now U.S. Patent No. 5,948,413. This application also claims priority to provisional application Serial No. 60/245,936, filed November 3, 2000.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

Not Applicable.

Reference to a "Computer Listing Appendix submitted on a Compact Disc"

Not Applicable.

BACKGROUND OF THE INVENTION

(1) Summary of the Invention

The present invention relates to antigen vaccines and methods of use thereof for the treatment of *Pythium insidiosum* infections, as well as prophylaxis against these infections, in humans and mammals. Further, the present invention relates to a method for preparing the preferred vaccine for the treatment which contains the intracellular and extracellular antigens of *Pythium insidiosum*. The present invention further relates to a mammal model for evaluating *Pythium insidiosum* vaccines and a method for monitoring the Th1 and Th2 response of a mammal in response to *Pythium insidiosum* vaccines.

(2) Description of Related Art

Infections caused by fungal and parafungal organisms are occurring with increasing frequency in patients with debilitating illnesses such as leukemia and AIDS, as well as those undergoing immunosuppressive therapy. Within this group of organisms are the traditional pathogenic fungi and a long list of newly recognized emerging opportunistic fungal and parafungal organisms. Among these emerging pathogens is the oomycete *Pythium insidiosum*, a fungal-like organism in the Kingdom Kromista, Phylum Pseudofungi. *Pythium insidiosum* is not only physiologically distinct from members of the Kingdom Fungi, but also differs physiologically. This may explain why anti-fungal drugs do not have any effect on pythiosis.

Pythiosis insidiosum particularly occurs in humans and lower animals in the tropical, subtropical, and temperate areas of the world (de Cock, W.A.W., et al., J. Clin. Microbiol. 25: 344-349 (1987)). The disease was first described in the beginning of the 20th century in equids of tropical and subtropical countries such as India and Indonesia as well as the USA. Soon, however, it was evident that the disease not only affected equids but other mammalian species such as humans. In lower animals, infections of the cutaneous tissues, lymphatic vessels, intestines, lungs, and bones have been found. In humans, a deadly arteritis infection, subcutaneous invasion, and keratitis occurs.

The drugs currently available to treat fungal infections have had little or no effect on *Pythium insidiosum*. Reports of treatment with either amphotericin B or surgery, which are commonly used to treat this disease in humans and lower animals, have indicated that 60% of the patients died of their infections. In cases of arterial invasion in humans, amphotericin B did not eliminate the infection (Rinaldi,

M.G., et al., Mycol. Obser. 9:7 (1989); and Thianprasit, M., Trop. Dermatol. 4: 1-4 (1990)), whereas in surgery the main problem has been to determine how much of the infected tissues has to be removed. Thus, relapses are common in surgically treated patients, who must also endure the pain and distress that such an invasive traumatic procedure inflicts on them.

The curative properties of *Pythium insidiosum* antigens was first noticed when some Costa Rican horses with pythiosis, which had been injected with *Pythium insidiosum* antigens in a skin test, were cured (Mendoza, L., et al., Equine pythiosis in Costa Rica: report of 39 cases. Mycopathologia 94: 123-126 (1986)). Simultaneously, another vaccine with similar curative properties was successfully used in horses with the disease in Australia (Miller, R. I., Aust. Vet. J. 57:3 77-382 (1981)). These two vaccines have been referred to in the literature as the Mendoza vaccine and the Miller vaccine, respectively (Newton, J. C., et al., The Compendium 15: 491-493 (1993)). Early reports indicated that the antigens used in the *Pythium insidiosum*-vaccine possessed unique characteristics, somewhat similar to the features of those reported in *Trichophyton verrucosum* (Gudding R., et al., Can. Vet. J. 36: 302-306 (1994)) and other immunotherapeutic vaccines (Foster, J. S., et al., Vet. Med. Small Ani. Clin. 71: 920 (1976); Pier, A. C., et al., Equine Pract. 15: 23-27 (1993)).

The Miller vaccine is prepared from sonicated hyphal antigens (Miller, R. E., Aust. Vet. J. 57: 377-382 (1981)), while the Mendoza vaccine is prepared from culture filtrate antigens (Mendoza, L., et al., Mycopathologia 94: 123-129 (1986)). Both the Miller vaccine and the Mendoza vaccine have been used to cure early cases of equine pythiosis, i.e., horses with pythiosis-caused lesions of 0.5 months or less in duration; however, neither vaccine can cure horses that are chronically infected with *Pythium insidiosum*, i.e.,

horses with pythiosis-caused lesions two or more months old (Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)). While both vaccines have cure rates of about 53% for early cases of pythiosis, the Mendoza vaccine has a longer shelf-life and milder side effects than the Miller vaccine (Miller, R. I., et al., J. Am. Vet. Med. Assoc. 182: 1227-1229 (1983)). The Mendoza vaccine, in addition to its immunotherapeutic properties, also showed some degree of protection against disease caused by *Pythium insidiosum*. However, this protection was later found to be of short duration (Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)).

In over 15 years of use, the Mendoza vaccine has been shown to be safe and consistently efficacious, curing more than 300 horses with pythiosis-caused lesions of short duration. However, the Mendoza vaccine can only cure early equine pythiosis, not chronic cases of this disease (Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)). Aside from the fact that the Mendoza vaccine can only cure early equine pythiosis cases, nothing was known about the immunogens involved in its curative properties nor the immune mechanisms that triggered the killing of *Pythium insidiosum*'s hyphae infected tissues.

In a recent study using SDS-PAGE and Western blot analysis, the presence of three immunodominant hyphal proteins were identified (Mendoza, L., et al., J. Clin. Microbiol. 30: 2980-2983 (1992)). While the Western blots revealed that the IgG in sera from horses with active pythiosis recognized most of the proteins of *Pythium insidiosum*, the Western blots also showed that three protein bands of the 32 kDa, 30 kDa, and 28 kDa were particularly prominent. More significant was the finding that antibodies against these three proteins persisted for long periods of time in the sera from successfully cured horses.

Even though there are two vaccines which can

be used to treat pythiosis, there remains a need for a vaccine which can cure pythiosis and prevent infection by *Pythium insidiosum*. In particular, there is a need for a vaccine that can be used to treat and cure patients who are in a chronic stage of the disease.

SUMMARY OF THE INVENTION

The present invention provides antigen vaccines and methods of use thereof for the treatment of *Pythium insidiosum* infections, as well as prophylaxis against these infections, in humans and mammals. Further, the present invention provides a method for preparing the preferred vaccine for the treatment which contains the intracellular and extracellular antigens of *Pythium insidiosum*. The present invention further provides a mammal model for evaluating *Pythium insidiosum* vaccines and a method for monitoring the Th1 and Th2 response of a mammal in response to *Pythium insidiosum* vaccines.

Therefore, the present invention provides a method for treatment of pythiosis in a human patient having pythiosis or prophylaxis against pythiosis which comprises (a) providing a vaccine consisting of intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum* and extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum* in a sterile aqueous solution; and (b) vaccinating the patient with the vaccine.

In further embodiment of the method, the vaccination is subcutaneous. Further still, the patient after vaccination is monitored for a change in a Th1 response and a Th2 response, wherein an increase in the Th1 response and a decrease in the Th2 indicates the patient has developed the Th1 response to the vaccine.

Further still, the present invention provides a method for the treatment of pythiosis in a mammal having pythiosis or prophylaxis against pythiosis in a

mammal which comprises (a) providing an injectable vaccine which comprises in a sterile aqueous solution in admixture (i) intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum*; and (ii) extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum*; and (b) vaccinating the mammal with the vaccine. Further still, the mammal after vaccination is monitored for a change in a Th1 response and a Th2 response, wherein an increase in the Th1 response and a decrease in the Th2 indicates the patient has developed the Th1 response to the vaccine.

In a further embodiment of the method, the antigens have been provided by (a) growing cells of the *Pythium insidiosum* in a culture medium and then (i) killing the cells; (ii) separating the killed cells from the culture medium so as to produce a first supernatant comprising the extracellular antigens secreted into the medium; and (ii) disrupting the cells in water to provide the intracellular cytoplasmic antigens in a second supernatant which is separated from the disrupted cells; and (b) separating the extracellular antigens from the first supernatant.

The present invention further provides a method for providing an injectable vaccine for treatment of pythiosis or prophylaxis against pythiosis which comprises (a) growing cells of *Pythium insidiosum* in a culture medium; (b) separating the cells from a first supernatant of the culture medium which contains extracellular proteins; (c) killing the cells; (d) disrupting the cells in sterile water; (e) separating the disrupted cells from the water to produce a second supernatant containing intracellular proteins; (f) mixing the first supernatant of step (b) with the second supernatant of step (e); (g) separating the combined proteins from the mixture of step (f); (h) mixing the separated proteins in sterile distilled water; and (i)

dialyzing the mixture of step (h) to remove low molecular weight components less than 10,000 MW to produce the vaccine.

5 The present invention further provides a method of testing a response in a mammal to a *Pythium insidiosum* vaccine including a cell-derived vaccine which comprises monitoring a Th1 response and a Th2 response of the mammal to the vaccine, wherein in mammals which are responding to the vaccine the Th1
10 response increases and the Th2 response decreases.

In a further embodiment of the method, the vaccine comprises antigens selected from the group consisting of intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum*,
15 extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum*, and combination of both.

In a further embodiment of the method, the antigens have been provided by growing cells of the *Pythium insidiosum* in a culture medium and then killing the cells, separating the killed cells from the culture medium so as to produce a first supernatant comprising the extracellular antigens secreted into the medium, separating the extracellular antigens from the first
20 supernatant, and disrupting the cells in water to provide the intracellular cytoplasmic antigens in a second supernatant which is separated from the disrupted cells.
25

In a further embodiment of the method, the mammal is infected with the *Pythium insidiosum* and the vaccine is for immunotherapy or the mammal is not infected with the *Pythium insidiosum* and the vaccine is for prophylaxis.
30

The present invention further provides a mammal model for testing a *Pythium insidiosum* vaccine including a cell-derived vaccine which comprises monitoring a Th1 response and a Th2 response of the
35

mammal in the mammal model to the vaccine, wherein in mammals which are responding to the vaccine the Th1 response increases and the Th2 response decreases.

5 In a further embodiment of the mammal model, the vaccine comprises antigens selected from the group consisting of intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum*, extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum*, and
10 combination of both.

In a further embodiment of the mammal model, the antigens have been provided by growing cells of the *Pythium insidiosum* in a culture medium and then killing the cells, separating the killed cells from the culture medium so as to produce a first supernatant comprising
5 the extracellular antigens secreted into the medium, separating the extracellular antigens from the first supernatant, and disrupting the cells in water to provide the intracellular cytoplasmic antigens in a second supernatant which is separated from the disrupted
10 cells. Preferably, the mammal in the mammal model is a rabbit.

In a further embodiment of the mammal model, the mammal is infected with the *Pythium insidiosum* and the vaccine is for immunotherapy or the mammal is
15 not infected with the *Pythium insidiosum* and the vaccine is for prophylaxis.

In a further embodiment of any one of the above embodiments of the present invention, the cells have been disrupted by sonication. In further
20 particular embodiments, the *Pythium insidiosum* is deposited as ATCC 74446; the culture medium is Sabouraud dextrose broth; the cells are killed with thimerosal; the disrupted cells are separated from the culture medium for the cells by centrifugation; the
25 intracellular cytoplasmic antigens in the second supernatant and the extracellular antigens in the first

supernatant are mixed to provide a mixture of antigens, precipitating the mixture of antigens with acetone to provide a precipitate, dissolving the precipitate in sterile distilled water to provide a solution of the antigens, and dialyzing the solution of antigens in sterile distilled water to remove low molecular weight components less than 10,000 MW to provide the vaccine; or, any combination thereof.

Further still, in any one of the above embodiments, the present invention provides the vaccines as disclosed in U.S. Patent Nos. 5,948,413 and 6,287,573, both to Mendoza. Briefly, the injectable vaccine for treatment of pythiosis or prophylaxis against pythiosis which comprises in a sterile aqueous solution an admixture of (a) intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum*; and (b) extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum*.

Preferably, the antigens in the injectable vaccine have been provided by (a) growing cells of the *Pythium insidiosum* in a culture medium and then (i) killing the cells; (ii) separating the killed cells from the culture medium so as to produce a first supernatant comprising the extracellular antigens secreted into the medium; and (ii) disrupting the cells in water to provide the intracellular cytoplasmic antigens in a second supernatant which is separated from the disrupted cells; and (b) separating the extracellular antigens from the first supernatant.

It is further preferable that the cells have been disrupted by sonication. Further still, it is preferable that the *Pythium insidiosum* is deposited as ATCC 74446. Further still, it is preferable that the culture medium is Sabouraud dextrose broth. Further still, it is preferable that the cells are killed with thimerosal. Further still, it is preferable that the

disrupted cells are separated from the culture medium by centrifugation. Further still, it is preferable that the intracellular cytoplasmic antigens in the second supernatant and the extracellular antigens in the first supernatant are mixed to provide a mixture of antigens, precipitating the mixture of antigens with acetone to provide a precipitate, dissolving the precipitate in sterile distilled water to provide a solution of the antigens, and dialyzing the solution of antigens in sterile distilled water to remove low molecular weight components less than 10,000 MW to provide the vaccine.

OBJECTS

It is therefore an object of the present invention to provide a method for treating pythiosis and for prophylaxis against pythiosis in humans and lower animals.

Further, it is an object of the present invention to provide vaccine compositions and methods for the preparation thereof.

Further still, it is an object of the present invention to provide a method for treating pythiosis or prophylaxis against pythiosis which is economical, reliable, and effective.

Further still, it is an object of the present invention to provide a method for monitoring the response of an animal to immunotherapy with the vaccine of the present invention.

Further still, it is an object of the present invention to provide a model system for testing the immune responses to the vaccine of the present invention.

These and other objects of the present invention will become increasingly apparent with reference to the following drawings and preferred embodiments.

DESCRIPTION OF THE DRAWINGS

Figure 1 shows microphotographs of the histopathological features of serial biopsies taken during successful immunotherapy using PIV in a horse with pythiosis.

Microphotograph A depicts the typical inflammatory reaction with eosinophils (arrows) degranulating around *Pythium insidiosum*'s hyphae and the Hoeppli-Splendore phenomenon that eventually increases in size, by the addition of more eosinophils, and it becomes the stony-like masses known as "kunkers" (H&E).

Microphotograph D shows typical silver stained hyphae of *Pythium insidiosum* from microphotograph A.

Microphotograph B shows few eosinophils (arrows) and numerous cytotoxic T lymphocytes and macrophages in biopsies taken seven days after successful immunotherapy. Kunkers were not found in biopsies from panel B (H&E).

Microphotograph E shows a silver-stained section of microphotograph B. Few hyphae were detected in the silver stained biopsies from microphotograph B. Most of them were observed as empty hyphal filaments (arrows).

Microphotograph C shows a predominantly mononuclear infiltrated 10 days after successful immunotherapy. The kunkers observed in microphotograph A, were also absent at this stage.

Microphotograph F shows a silver-stained section of microphotograph C. No hyphae of *Pythium insidiosum* were found in the biopsies.

Figure 2 illustrates a working hypothesis based on data from cured humans and equines which explains the therapeutic properties of PIV.

DESCRIPTION OF PREFERRED EMBODIMENTS

All patents, patent applications, government publications, government regulations, and literature

references cited in this specification are hereby incorporated herein by reference in their entirety. In case of conflict, the present description, including definitions, will control.

5 The term "extracellular antigens" refers to antigens or proteins which are secreted from *Pythium insidiosum* during its *in vitro* cultivation. These are antigens during the life of *Pythium insidiosum* which are excreted or secreted from the organism.

10 The term "intracellular antigen" refers to cytoplasmic antigens or proteins which are in the cytoplasm of *Pythium insidiosum* during its *in vitro* cultivation. These are antigens which during the life of *Pythium insidiosum* do not appear outside the organism.

15 The term "intracellular cytoplasmic antigen" refers to intracellular antigens found in the cytoplasm.

20 The terms "antigens" and "proteins" are used interchangeably. Therefore, unless otherwise stated, the use of one term does not exclude the other term.

25 As set forth herein and in U.S. patent Nos. 5,948,413 and 6,287,573, both to Mendoza, the present invention relates to an injectable vaccine for treatment of pythiosis which comprises in sterile aqueous solution an admixture of (a) intracellular proteins separated from disrupted cells of *Pythium insidiosum*; and (b) extracellular proteins from a supernatant from growing the cells of the *Pythium insidiosum*.

30 The present invention also relates to a method for providing an injectable vaccine for treatment of Pythiosis which comprises (a) growing cells of *Pythium insidiosum* in a culture medium; (b) separating the cells from a first supernatant of the culture medium which contains extracellular proteins; (c) killing the cells;
35 (d) disrupting the cells in sterile water; (e) separating the disrupted cells from the water to produce a second supernatant containing intracellular proteins;

(f) mixing the first supernatant of step (b) with the second supernatant of step (e); (g) separating the combined proteins from the mixture of step (f); (h) mixing the separated proteins in sterile distilled water; and (i) dialyzing the mixture of step (h) to remove low molecular weight components less than 10,000 MW to produce the vaccine.

The present invention further relates to a method for the treatment of Pythiosis in a mammal having the disease which comprises (a) providing an injectable vaccine which comprises in a sterile aqueous solution in admixture: (1) an intracellular proteins separated from disrupted cells of *Pythium insidiosum*; and (2) extracellular proteins from a supernatant from growing the cells of the *Pythium insidiosum*; and (b) vaccinating the mammal with the vaccine.

Further still, the present invention relates to a method for treatment of pythiosis in human patients having the disease which comprises (a) providing a vaccine containing separated proteins of *Pythium insidiosum* in a sterile aqueous solution; and (b) vaccinating the patient with the vaccine.

Therefore, the present invention provides method for using an injectable *Pythium insidiosum* vaccine (PIV) for use as a therapeutic vaccine to cure pythiosis or as a prophylactic vaccine against pythiosis. The PIV comprises in a sterile aqueous solution an admixture of intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum* and extracellular antigens secreted into the medium of the medium used to grow the cells of the *Pythium insidiosum*. The intracellular cytoplasmic antigens include the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens as determined by SDS polyacrylamide gel electrophoresis.

Preferably, the PIV contains at least 2 to 3 mg/ml of the extracellular and intracellular antigens.

The preferred vaccination dosage for the PIV is at least about 1 to 2 mg/kg body weight of the vaccinee.

5 The route of administration for the PIV of the present invention includes, but is not limited to, intramuscular, intraperitoneal, intradermal, subcutaneous, intravenous, intra-arterial, intra-ocular, and trans-dermal or by inhalation or suppository. The preferred routes of administration include intramuscular, intraperitoneal, intradermal, and subcutaneous injection. Most preferably, the PIV is injected intramuscularly. The PIV can be administered by means including, but not limited to, syringes, needle-less injection devices, or microprojectile bombardment gene guns.

15 The PIV of the present invention is formulated in a pharmaceutically accepted carrier according to the mode of administration to be used. In cases where intramuscular injection is preferred, a sterile water or isotonic formulation is preferred. Generally, additives for isotonicity can include sodium chloride, dextrose, mannitol, sorbitol, and lactose. In particular cases, isotonic solutions such as phosphate buffered saline are preferred. The formulations can further provide stabilizers such as gelatin and albumin. In some embodiments, a vaso-constriction agent is added to the formulation. An adjuvant for the vaccine is EMULSIGEN (MVP Labs, Ralston, Nebraska), which is a paraffin oil in a water emulsion, which can be used in food animals. Freund's Incomplete Adjuvant, which is 15 percent by weight mannide monooleate and 85% paraffin oil, available from Difco, Detroit, Michigan, can be used in non-food (i.e. laboratory animals). The adjuvants aid in slowly releasing the vaccine into the animal and can potentiate the immune response. Any commercial oil emulsion adjuvants can be used such as vitamin E. The most preferred carrier is sterile water or an aqueous saline solution, particularly when the vaccinee is a

human.

5 The pharmaceutical preparation according to
the present invention are provided sterile and pyrogen
free. However, it is well known by those skilled in the
art that the preferred formulations for the
pharmaceutically accepted carrier which comprise the PIV
of the present invention are those pharmaceutical
carriers approved in the regulations promulgated by the
United States Department of Agriculture, or equivalent
10 government agency in a foreign country such as Canada or
Mexico, for vaccines intended for veterinary
applications. Therefore, a pharmaceutically accepted
carrier for commercial production of the PIV of the
present invention is a carrier that is already approved
or will at some future date be approved by the
15 appropriate government agency in the United States of
America or foreign country.

Inoculation of the vaccinee with the PIV is
preferably by a single vaccination. In another
20 embodiment of the present invention, the vaccinee is
subjected to a series of vaccinations to produce a full,
broad immune response. When the vaccinations are
provided in a series, the vaccinations can be provided
between about 24 hours apart to two weeks or longer
25 between vaccinations. In certain embodiments, the
vaccinee is vaccinated at different sites
simultaneously.

A preferred *Pythium insidiosum* for preparing
the PIV was deposited with the American Type Culture
Collection under the Budapest Treaty as ATCC 74446. It
30 is available upon request by name and number. All
restrictions on distribution of ATCC 74446 are
irrevocably removed on granting of a patent on this
application. The address of the American Type Culture
Collection is 10801 University Boulevard, Manassas,
35 Virginia 20110-2700.

In further embodiments of the present

invention, the PIV can be combined with immunizing or non-immunizing components for other diseases to produce a multivalent vaccine or with other medicaments, particularly antibiotics. The antibiotics can be used prior to vaccination.

In the following Example 1, the PIV of the present invention was prepared by adding cytoplasmic antigens to the earlier *Pythium insidiosum*-vaccine (Mendoza vaccine; Mendoza et al., Mycopathologica 119: 89-95 (1992)). In Example 2, the PIV of Example 1 was tested in horses with chronic pythiosis insidiosi, only 48% of the horses were cured. All horses with acute pythiosis insidiosi were cured with this new vaccine. One advantage of the PIV is that the earlier Mendoza vaccine always failed in chronic cases. Example 3 shows preparation of monoclonal antibodies against the immunodominant antigens or proteins comprising the PIV. Example 4 shows the production of the immunodominant antigens by recombinant DNA methods. In Example 5, the PIV was successfully tested in a Thai boy with pythiosis insidiosi. This Thai patient had been diagnosed with an infection caused by *Pythium insidiosum* in his external carotid artery. In spite of efforts to treat the infection with traditional methods, the patient did not show improvement. As a last resort PIV was given to him. The patient was clinically cured.

Traditionally, vaccines have been used primarily for prophylactic purposes. The use of vaccines for the treatment of ongoing diseases, even though an old idea, has only recently received attention (Cohen, J., Science 264: 503-505 (1994)). The long-held medical dogma that vaccines are only for prevention has been challenged by scientists working toward the development of immunotherapeutic vaccines against viruses (Burke, D. S., Vaccine 11: 883-890 (1993)), parasites (Convit, J., et al., Trans. Royal Soc. Trop. Med. Hyg. 87: 444-448 (1993)), bacteria (Standford, J.

L., Trop. Geograph. Med. 46: 93-107 (1994)), fungal (Gudding, R., et al., Can. Vet. J. 36: 302-306 (1995)), and parafungal pathogens (Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)).

5 Despite impressive data originated by the PIV and other curative vaccines, strong skepticism still exists against the use of therapeutic vaccines as weapons for the treatment of infectious diseases. The skeptics have argued that when a host is invaded by an
10 organism, its immune system will mount an immune response that will eventually eliminate the invader. They argue, however, that if the immune system fails, the only treatment left is the use of drugs. However, the findings presented herein using the PIV and data
15 elsewhere using other therapeutic vaccines have indicated that a new line of research is necessary to investigate the mechanism by which these therapeutic vaccines elicit an immunological reaction that kills the pathogens in infected tissues.

20 The mechanisms underlying the response to PIV are not well understood. However, based on histopathological and immunological studies in cured horses, it was found that the cellular immune response plays a major role in the clearance of *Pythium*
25 *insidiosum* from infected tissues (Mendoza, L., et al., Mycopathologia 94: 123-129 (1986); Miller, R. I., Aust. Vet. J. 67: 377-382 (1981); Newton, J. C., et al., Compendium 15:491-493 (1993); and Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)). These studies have
30 all shown that after successful immunotherapy, the eosinophilic inflammatory reaction, typical of this disease, gradually changed to a mononuclear immunoresponse. Numerous macrophages, lymphocytes (cytotoxic), and plasma cells had replaced the
35 eosinophilic granuloma. Surprisingly, the mononuclear cells surrounded and killed *Pythium insidiosum's* hyphae, eliminating the pathogen from the affected tissues.

This observation has been corroborated by the failure to recover *Pythium insidiosum* from the tissue of equines cured by immunotherapy (Newton, J. C., et al., Compendium 15: 491-493 (1993) and Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)).

Based on the PIV data accumulated in the past 15 years in equine pythiosis, it is strongly believed that the *Pythium insidiosum* vaccine displays to the host's immune system epitopes that are not well presented during natural infection. This scenario is possible since *Pythium insidiosum*'s hyphae are always sequestered inside eosinophilic granulomas. Thus, *Pythium insidiosum* is probably using the degranulated eosinophils to hide important epitopes from the host's immune system.

Example 6 provides evidence that the mechanism by which the PIV of the present invention provides its immunotherapeutic effect is by down regulating the T helper 2 (Th2) subset and activating the T helper 1 (Th1) subset. In the natural infection, the hyphae produce extracellular antigens which direct the host's immune system to mount a Th2 response which stimulates production of IgE, IgM, and IgG. The IgE triggers the production of eosinophils and mast cells which degranulate over the hyphae thereby protecting the hyphae which enables the *Pythium insidiosum* to multiply and produce overwhelming quantities of extracellular antigens which lock the host's immune response in the Th2 mode. The PIV, by switching the immune response from a Th2 response to a Th1 response, triggers a mononuclear cell mediated immune (CMI) response in the host which is composed mostly of cytotoxic T lymphocytes (CTL) and macrophages. The CTLs and macrophages damage and destroy the hyphae of *Pythium insidiosum* which prevents the production of the extracellular antigens that are required to maintain the Th2 subset. Because in the absence of the extracellular antigens the host's

immune system is switched to and remains in the Th1 mode, the host's CTLs and macrophages are able to destroy the *Pythium insidiosum* thereby curing the host of the *Pythium insidiosum*. Furthermore, the results indicate that the PIV of the present invention can be used for prophylaxis as well because the PIV stimulates a Th1 response which then attacks and destroys the hyphae as they are formed. As a consequence, the *Pythium insidiosum* cannot establish an infection.

Example 7 provides a mammal model for evaluating *Pythium insidiosum* vaccines. The mammal model for testing a *Pythium insidiosum* vaccine including cell-derived vaccine comprises monitoring a Th1 response and a Th2 response of the mammal in the mammal model to the vaccine, wherein in mammals which are responding to the vaccine the Th1 response increases and the Th2 response decreases. The preferred mammal is a rabbit. Example 7 also provides a method for the testing a response in a mammal to a *Pythium insidiosum* vaccine including a cell-derived vaccine which comprises monitoring a Th1 response and a Th2 response of the mammal to the vaccine, wherein in mammals which are responding to the immunotherapy, the Th1 response increases and the Th2 response decreases.

The vaccine comprises antigens selected from the group consisting of intracellular cytoplasmic antigens separated from disrupted cells of *Pythium insidiosum*, extracellular antigens secreted into a medium for growing the cells of the *Pythium insidiosum*, and combination of both. The antigens can be produced *in vitro* by recombinant DNA technology, by DNA vaccines, or derived or isolated from the *Pythium insidiosum*.

When the *Pythium insidiosum* is the source of the antigens, the antigens are provided by growing the *Pythium insidiosum* in a culture medium and then killing it, separating the killed *Pythium insidiosum* from the culture medium so as to produce a first supernatant

comprising the extracellular antigens secreted into the medium, separating the extracellular antigens from the first supernatant, and disrupting the cells in water to provide the intracellular cytoplasmic antigens in a second supernatant which is separated from the disrupted cells.

The following examples are intended to promote a further understanding of the present invention.

EXAMPLE 1

This example shows the steps for preparing the PIV of the present invention.

1. *Pythium insidiosum* strain ATCC 74446, was transferred to a 1.0-liter flask containing 500 ml of Sabouraud dextrose broth (Difco, Detroit, Michigan).

2. Cultures were incubated at 37° C for five days on a rotary shaker rotating at 150 rpm.

3. Cultures were killed with Merthiolate (thimerosal) (0.02% wt/vol), filtered to separate the cells (hyphae) from the liquid phase containing the secreted extracellular antigens of *Pythium insidiosum*. The liquid phase was saved in a sterile container to be used in step 6.

4. The cell mass obtained in step 3, was washed twice with sterile distilled water and disrupted by sonication until 100% of the hyphae were fragmented. This step releases the intracellular cytoplasmic antigens. Other methods for disrupting the cells such as a French press can be used.

5. The disrupted cell mass obtained in step 4, was centrifuged at 5,000 x g for 20 minutes to separate the cell debris which pellet from the intracellular cytoplasmic antigens which remain in the supernatant fraction.

6. The supernatant fraction was decanted from the pellet and the pellet was discarded. Then the supernatant fraction was added to the liquid phase in

step 3 to make a mixture of the supernatant fraction and the liquid phase.

5 7. To confirm the presence of the immunodominant proteins in the mixture or the supernatant fraction before mixing, a sample of the mixture or supernatant fraction was subjected to SDS-PAGE electrophoresis and Western blot analysis to verify that the supernatant fraction contained the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens identified in
10 Mendoza et al. (J. Clin. Microbiol. 30: 2980-29-83 (1992)).

15 8. After visualization of the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens by Western blot analysis, the mixture was then precipitated with an equal volume of acetone and pelleted at 20,000 x g for 30 minutes in a refrigerated centrifuge to concentrate the intracellular cytoplasmic antigens and extracellular secreted antigens and remove acetone soluble material.

20 9. The supernatant fraction was decanted and the pellet was dissolved in sterile distilled water to make a solution with a final protein concentration of about 2.0 mg/ml.

25 10. The solution was dialyzed using a membrane cut off point of 10,000 MW to remove low molecular weight material to produce the PIV which contains the intracellular cytoplasmic antigens, which includes the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, and the extracellular secreted antigens.

30 11. The sterility of the PIV was confirmed by culturing 100 µl of the PIV on blood agar and Sabouraud dextrose broth.

35 12. The PIV was stored at 4° C or lyophilized until use. When the PIV was stored lyophilized, it was resuspended in sterile distilled water to a final protein concentration of 2.0 mg/ml before use.

In some cases, the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens were purified by preparative SDS

polyacrylamide gel electrophoresis. In such cases, following electrophoresis, the 32 kDa, 30 kDa, and 28 kDa antigens were cut from the acrylamide gels and purified. The 32 kDa, 30 kDa, and 28 kDa immunodominant antigens were mixed to make a mixture which a portion of which was added to the Mendoza vaccine to produce a modified Mendoza vaccine with a final protein concentration of about 2.0 µg/ml. A western blot analysis was then performed on the modified Mendoza vaccine to verify the presence of the 32 kDa, 30 kDa, and 28 kDa antigens. The modified Mendoza vaccine was stored at 4° C or lyophilized until use. When the modified Mendoza vaccine was stored lyophilized, it was resuspended in sterile distilled water to a final protein concentration of 2.0 mg/ml before use.

EXAMPLE 2

One major drawback in evaluating the PIV or any other vaccine against *Pythium insidiosum* is the lack of an animal model. The only animal in which the disease can be successfully reproduced is the rabbit (*Oryctolagus cuniculus*). However, no systematic studies have been conducted to evaluate the effectiveness of the rabbit as an experimental model. Therefore, evaluations of the PIV has been carried out only in horses with the disease. The diagnosis of pythiosis in the treated equines was verified either by serology, by culture, by histopathology, or by all three methods. Based on the fact that neither the Miller vaccine (a cell-mass vaccine consisting solely of disrupted *Pythium insidiosum* cell debris) nor the Mendoza vaccine (a soluble concentrated antigen vaccine consisting solely of *Pythium insidiosum* extracellular antigens) cured infected horses after 60 days or more of infection, seven horses were selected with chronic pythiosis (greater than 60 days after infection, some more than 100 days after infection) and three horses

with acute pythiosis (less than 60 days after infection), to conduct a vaccination trial with the PIV as prepared as in Example 1.

5 The results indicated that the efficacy of the PIV, which contains both cytoplasmic antigens, which included the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, and extracellular antigens, was remarkably superior to the Mendoza vaccine which contained only extracellular antigens. Of the seven horses with
10 chronic pythiosis vaccinated four were cured, two did not respond, and one initially responded but died later. All of the cured horses developed a mild inflammatory reaction at their vaccination sites. However, the three horses that did not respond to the vaccinations did not
15 develop such a reaction. Those horses had had their infections for more than 100 days and were considered to be anergic. The PIV also cured all of the early cases of pythiosis.

20 The results suggest that 1) the presence of the cytoplasmic antigens, which included the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, directly enhanced the efficacy of the Mendoza vaccine which always failed in chronic cases (>60 days) (Mendoza, L., et al., Mycopathologia 119: 89-95 (1992)); 2) the
25 cytoplasmic antigens are directly affect the immunotherapeutic properties of the Mendoza vaccine, and 3) the cytoplasmic antigens play a role in the immunology of *Pythium insidiosum* infection.

30 The findings also confirmed that the response to vaccination is directly related to the immune status of the infected horse. Although the PIV's main attribute is its therapeutic ability to cure chronic equine pythiosis cases, the extracellular antigens in the PIV allowed the PIV to have all of the properties of
35 the Mendoza vaccine. These include, production of a mild inflammatory reaction at the site of vaccination in cured, but not in unresponsive equines, and a 100% cure

rate in early cases. The cure rate using the Mendoza vaccine was 48%. After addition of the cytoplasmic antigens, which included the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, the cure rate increased to 70%.
5 The enhancement in efficacy was directly related to the addition of the cytoplasmic antigens, which included the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, to the extracellular antigens.

10 EXAMPLE 3

This example shows the preparation of monoclonal antibodies that recognize the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens.

15 The 32 kDa, 30 kDa, and 28 kDa immunodominant antigens are separated on SDS gels as taught in Mendoza et al (J. Clin. Microbiol. 30: 2980-29-83 (1992)). Following electrophoresis, the prominent proteins are cut from the acrylamide gels and purified using standard protein purification methods. Then the purified
20 antigens are used to make monoclonal antibodies according to the methods in *Antibodies, A Laboratory Manual*, eds. Harlow and Lane, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988), well known to those skilled in the art as a source for
25 methods for making polyclonal and monoclonal antibodies.

BALB/c mice are immunized with an initial injection of 1.0 µg of the 32 kDa, 30 kDa, or 28 kDa immunodominant antigens per mouse mixed 1:1 with Freund's complete adjuvant. After two weeks, a booster
30 injection of 1.0 µg of the antigen is injected into each mouse intravenously without adjuvant. Three days after the booster injection the mouse serum is checked for antibodies to the 32 kDa, 30 kDa, or 28 kDa immunodominant antigen. If positive, a fusion is
35 performed with a mouse myeloma cell line. Mid log phase myeloma cells are harvested on the day of fusion, checked for viability, and separated from the culture

medium by low-speed centrifugation. Then the cells are resuspended in serum-free Dulbecco's Modified Eagle's medium (DMEM).

5 The spleens are removed from the immunized mice and washed three times with serum-free DMEM and placed in a sterile Petri dish containing 20 ml of DMEM containing 20% fetal bovine serum, 1 mM pyruvate, 100 units penicillin, and 100 units streptomycin. The cells are released by perfusion with a 23 gauge needle. 10 Afterwards, the cells are pelleted by low-speed centrifugation and the cell pellet is resuspended in 5 ml 0.17 M ammonium chloride and placed on ice for several minutes. Then 5 ml of 20% bovine fetal serum is added and the cells pelleted by low-speed 15 centrifugation. Afterwards, the cells are resuspended in 10 ml DMEM and mixed with myeloma cells to give a ratio of 3:1. The cell mixture is pelleted by low-speed centrifugation, the supernatant fraction removed, and the pellet allowed to stand for 5 minutes. Next, over 20 a period of 1 minute, 1 ml of 50% polyethylene glycol (PEG) in 0.01 M HEPES pH 8.1 at 37°C is added. After 1 minute incubation at 37°C, 1 ml of DMEM is added for a period of another 1 minute, then a third addition of DMEM is added for a further period of 1 minute. 25 Finally, 10 ml of DMEM is added over a period of 2 minutes. Afterwards, the cells are pelleted by low-speed centrifugation and the pellet resuspended in DMEM containing 20% fetal bovine serum, 0.016 mM thymidine, 0.1 hypoxanthine, 0.5 μ M aminopterin, and 10% hybridoma cloning factor (HAT medium). The cells are then plated 30 into 96-well plates.

After 3, 5, and 7 days half the medium in the plates is removed and replaced with fresh HAT medium. After 11 days, the hybridoma cell supernatant is 35 screened by an ELISA assay. In this assay, 96-well plates are coated with the appropriate 32 kDa, 30 kDa, or 28 kDa immunodominant antigen. One hundred μ l of

supernatant from each well is added to a corresponding well on a screening plate and incubated for 1 hour at room temperature. After incubation, each well is washed three times with water and 100 μ l of a horseradish peroxide conjugate of goat anti-mouse IgG (H+L), A, M (1:1,500 dilution) is added to each well and incubated for 1 hour at room temperature. Afterwards, the wells are washed three times with water and the substrate OPD/hydrogen peroxide is added and the reaction is allowed to proceed for about 15 minutes at room temperature. Then 100 μ l of 1 M HCl is added to stop the reaction and the absorbance of the wells is measured at 490 nm. Cultures that have an absorbance greater than the control wells are removed to 2 cm² culture dishes, with the addition of normal mouse spleen cells in HAT medium. After a further three days, the cultures are re-screened as above and those that are positive are cloned by limiting dilution. The cells in each 2 cm² culture are counted and the cell concentration adjusted to 1 x 10⁵ cells per ml. The cells are diluted in complete medium and normal mouse spleen cells are added. The cells are plated in 96-well plates for each dilution. After 10 days, the cells are screened for growth. The growth positive wells are screened for antibody production; those testing positive are expanded to 2 cm² cultures and provided with normal mouse spleen cells. This cloning procedure is repeated until stable antibody producing hybridomas are obtained. Then the identified stable hybridomas are progressively expanded to larger culture dishes to provide stocks of the cells.

Production of ascites fluid is performed by injecting intraperitoneally 0.5 ml of pristane into female mice to prime the mice for ascites production. After 10 to 60 days, 4.5 x 10⁶ cells are injected intraperitoneally into each mouse and ascites fluid is harvested between 7 and 14 days later.

An alternate method for screening hybridomas

for antibody production is as follows. *Pythium insidiosum* is heat-denatured in 0.5 M Tris (pH 7.4) with 10% SDS, 20% glycerol and 5% 2-mercaptoethanol. The denatured antigens are separated by SDS-polyacrylamide gel electrophoresis in a 12-20% (v/v) linear gradient gel with a 4% (v/v) stacking gel. The separated antigens are electrophoretically transferred to Western PVDF membranes at 100 volts for 1.5 hours, then 150 volts for 0.5 hours. The membranes are then blocked overnight in 1% by volume bovine serum albumen in 0.5% Tween-Tris buffered saline (Blocking buffer). The blots are air-dried and stored frozen. Prior to use, the membranes are incubated with bovine serum albumin in Blocking buffer at a range of 1:10 to 1:100 ratio for two hours. Afterwards, the membranes are washed in 0.5% Tween-Tris buffered saline and then incubated with monoclonal antibodies from the various hybridoma clones. The membranes are developed as disclosed in the prior art, e.g., Granstrom et al., J. Vet. Diag. Invest. 5: 88-90 (1993) or *Antibodies, A Laboratory Manual*, eds. Harlow and Lane, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988).

Hybridomas that successfully produce monoclonal antibodies against various epitopes of the 32 kDa, 30 kDa, or 28 kDa immunodominant antigen are expanded as above, and used to make monoclonal antibodies for the antigen-based immunoassay and for identifying cDNA library clones in Example 4 that contain *Pythium insidiosum* DNA which express either the 32 kDa, 30 kDa, or 28 kDa immunodominant antigen.

EXAMPLE 4

This example shows the preparation of a cDNA library that expresses the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens. The methods for making and screening cDNA expression libraries are well known to those skilled in the art and are described in *Molecular*

Cloning: A Laboratory Manual, Second Edition, edited by Sambrook et al. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). The monoclonal antibodies made as in Example 3 are used to screen the library for clones that express the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens.

Clones encoding the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens can be used to dissect, at a molecular level, the components that provide the antigens their protective and curative properties. The genes can be used to express the immunodominant antigens in an expression vector in *E. coli* and the expressed antigens purified and combined with the extracellular antigens to provide a vaccine that contains only the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens mixed with the extracellular antigens.

EXAMPLE 5

This Example shows the use of the PIV of Example 1 to successfully treat a Thai boy with a life threatening pythiosis insidiosus arteritis.

A 14 year-old boy was admitted to the Ramathibodi Hospital, Bangkok, Thailand, with a history of 10 days of progressive headache. The illness had begun 16 days before admission in November 1995. Previous to the symptoms, he had developed a small skin injury on the posterior portion of his neck while swimming in a flooded area near a rice field. Four days after the skin injury, he developed three acne-like nodules at the injured site. He was then admitted to a local hospital with a severe headache and soft tissue swelling at the occiput. The swollen mass returned to normal after two days of dexamethazone treatment. The patient, however, continued to have severe headaches and developed a left facial nerve palsy before admission to the Ramathibodi Hospital.

The boy had a history of post splenectomy β -

thalassemia hemoglobin E disease of four years duration. He had received at least three blood transfusions per year after his operation. Headache, bilateral facial nerve palsy, and progressively extensive facial cellulitis were recorded on admission. Empirical antibiotic treatment with cefotaxime 100 mg/kg/day and chloramphenicol 75 mg/kg/day were prescribed without success. A computerized tomography (CT) scan of the head and neck showed diffuse cellulitis. Abscesses in the bilateral retromolar fossa and in both ears were also observed. After surgical drainage of the abscesses, the pain and headache were relieved and the soft tissue swelling subsided. A non-sporulating fungus-like organism was isolated in pure culture from tissue taken from the left and the right pinna. Because of the possibility of a fungal infection amphotericin B 0.5 mg/kg/day increasing to 1 mg/kg/day was administered. The isolate was later identified as *Pythium insidiosum*.

Although the abscess and cellulitis subsided, one week later, however, the pain and headache reappeared. Swelling of the left side of his tongue was also noticed. Saturated potassium iodide (1 g/ml) 3 ml/day that was increased gradually to 9 ml was prescribed. Despite this treatment, no clinical improvement was observed. Magnetic resonance images (MRI) of the head and neck demonstrated soft tissue involvement and regional lymph node enlargement. Surgical exploration of the left parapharynx and masseteric space was performed. During surgical exploration, the left abnormal cervical lymph nodes and the abnormal left great auricular nerve were removed. Histopathologically, the material showed follicular hyperplasia with sinus histiocytosis and granulomatous inflammation and aseptate hyphal elements of *Pythium insidiosum*. After failure with amphotericin B and iodides, chemotherapy with 300 mg/day of ketoconazole

was initiated. Granulocyte macrophage colony stimulating factor (GM-CSF) was given 5 days immediately post surgical exploration.

5 The headache and swollen tongue improved after surgical intervention. Although treatment with ketoconazole and iodides continued, pain and headache reappeared three weeks later. A CT angiogram revealed an aneurysm in the left external carotid artery 1.0 cm above the bifurcation and stenosis with irregular walls of the internal carotid artery. A third surgical intervention was performed on February 1, 1996 to remove the aneurysm. The excised tissue was oval in shape 2.5-4 cm in diameter with necrotic-like material within its lumen. Histopathologically, eosinophils, macrophages, 10 CD3 positive T-cells, plasma cells, and hyphal elements of *Pythium insidiosum* were observed within the lumen and the vessel's wall. Pain and headache disappeared immediately after the surgical intervention.

15 Five weeks after surgery, headache and swelling tissue returned. An MRI and an MRA of the neck revealed the persistence of cervical and paracervical lymph node enlargement and persistent stenosis of the left internal carotid artery. These findings suggested that *Pythium insidiosum* had invaded that artery as well. 20 Surgical removal of the left internal carotid artery was not recommended. Since amphotericin B, ketoconazole, iodides, surgery, and two courses of GM-CSF alone were ineffective in controlling the infection, the PIV was suggested as a last resort treatment. In particular, 25 because of the success of the PIV in animals with pythiosis insidiosum.

30 Vaccine administration. A dose of 100 µl of the 2 mg/ml PIV had been used as shown in Example 2 to vaccinate horses with the disease. In successfully 35 treated horses, an inflammatory reaction always developed at the site of vaccination. This inflammatory response indicated not only that the host's immune

system was functioning, but it also predicted that the horse probably would be cured by the vaccine. Anergic horses with proven pythiosis insidiosus never developed such a reaction to the vaccine and did not respond to the immunotherapy (Mendoza, L., et al., Mycopathologia 94: 123-129 (1986); Newton, J. C., et al., Equine pythiosis: An overview of immunotherapy. Compendium 15: 491-493 (1993); and Miller, R. I., et al., J. Am. Vet. Med. Assoc. 182: 1227-1229 (1983)).

To avoid an excessive immunoresponse in the young boy with *Pythium insidiosum* arteritis, several dilutions of the original PIV were tested before the trial started. One hundred µl of each PIV dilution (1:100 to 1:100,000) were injected as a skin test on his right forearm. A mild inflammatory reaction was observed only with the 1:100 dilution of the PIV. Thus, the undiluted batch of PIV was selected. One hundred µl of the 2 mg/ml PIV was subcutaneously injected in the patient's left shoulder.

Twenty hours after vaccination, a wheal and flare reaction had developed at the injection site. Forty-eight hours post vaccination, the wheal reaction attained its maximum size of 11 cm in diameter. No other side effects occurred except for itching at the vaccination site. The skin reaction disappeared 10 days post vaccination. Fourteen days after the first dose, the facial and tongue swelling had diminished. The same day a second vaccination was performed on the patient's right shoulder. Forty-eight hours later the wheal reaction at the vaccination had attained a diameter of eight centimeters. Two weeks after the second vaccination the patient's headache had disappeared, his facial and left tongue swelling were dramatically diminished, and the enlarged cervical lymph node had reduced in size. For the first time since his admission the patient's weight had increased by 4.0 kg four weeks post vaccination. The boy was considered clinically

cured one year after the first vaccination.

An MRI performed 6 weeks after the first vaccination, showed a decrease in the thickening of the soft tissue and less soft tissue enhancement of the left side of his tongue. An MRA of the neck released significant improvement of the stenosis of the proximal left internal carotid artery. The MRI and MRA twelve months post vaccination showed no infiltrations in the soft tissue and a normal left internal carotid artery.

A serum sample collected during the initial weeks post-admission gave a negative result in an ID for pythiosis. Although the ID test in equine pythiosis is a reliable test, some negative results have been reported in humans and dogs with proven pythiosis (Chetchotisakd, P., et al., J. Med. Assoc. Thailand 75: 248-254 (1992); and Wanachiwanawin, W., et al., Trans. Royal Soc. Trop. Med. Hyg. 87: 296-298 (1993)). When the serum sample was tested in a new *Pythium insidiosum*-ELISA, positive titers of 1:6,400 were recorded. To monitor the vaccination's progress, sera collected one, two, six and twelve months post-vaccination were also evaluated with the ELISA. A decrease in titers from 1:6,400 to 1:800 after 6 months post vaccination indicated that the *Pythium insidiosum* may have been eliminated from the infected tissues, a finding that substantiated the clinical data. Later serum samples showed that the antibody titer against *Pythium insidiosum* continued to decrease. However, as has been previously reported in equines cured by immunotherapy, low antibody titers may persist for years after the cure (Mendoza, L., et al., J. Clin. Microbiol. 30: 2980-2983 (1992)).

The response of the patient to the PIV vaccine was remarkable. Besides the wheal and flare reaction at the site of vaccination no deleterious side effects developed. Within four weeks after immunotherapy his headaches had disappeared, tissue swelling decreased,

and he gained 4.0 Kg in weight. Although the full strength vaccine was used (2 mg/ml) the patient tolerated the PIV very well. The success obtained with the immunotherapy in this particular case indicates that PIV can be used as an alternative therapy for human pythiosis insidiosus. This finding is of importance because the available antifungal drugs have little effect on this emerging pathogen. This is the first human pythiosis insidiosus arteritis case that has been treated and cured by the immunotherapeutic PIV.

EXAMPLE 6

The objective of this example was to assess the immunotherapeutic properties of the *Pythium insidiosum* vaccine (PIV) of the present invention in horses, dogs and a cat with pythiosis and to determine the immunological response in horses after successful immunotherapy.

This example provides evidence that down regulation of T helper 2 (Th2) and activation of T helper 1 (Th1) subsets may be responsible for the immunotherapeutic properties of the PIV of the present invention. The results suggest that PIV causes a shift in the immune system of the infected host from a Th2 response to a Th1 response. The cytoplasmic intracellular antigens, which include the 32 kDa, 30 kDa, and 28 kDa immunodominant antigens, in the PIV effect the down regulation of the Th2 response in hosts that are infected with pythiosis and activation of the Th1 response which enables the PIV to cure pythiosis. These results provide an explanation as to why PIV is efficacious both as a therapeutic vaccine for treatment of hosts infected with *Pythium insidiosum* and as a prophylactic vaccine to prevent *Pythium insidiosum* infection in a vaccinee.

Animals used in the example. Between 1996 and 1998, 19 horses, five dogs, and one cat diagnosed with

pythiosis by serological and other methods, were selected for immunotherapy with the PIV of the present invention. The injected animals were clinically and serologically evaluated for more than two months after immunotherapy. Histopathological studies using H&E and silver stains were performed on tissue samples from some of the animals. One horse from the cured group was selected for multiple biopsies.

Serological Assays to Diagnose Pythiosis. Two serological assays were used to diagnose pythiosis in the infected animals. These included immunodiffusion (ID) (Mendoza et al., J. Clin. Microbiol. 23: 813-816 (1986)) and ELISA (Mendoza et al., Clin. Diag. Lab. Immunol. 4: 715-718 (1997)). The antigen for ID was prepared as per Mendoza et al. (Mendoza et al., J. Clin. Microbiol. 23: 813-816 (1986); Mendoza et al., Clin. Diag. Lab. Immunol. 4: 715-718 (1997)). Briefly, *Pythium insidiosum* (ATCC 74446, type strain) was incubated in Sabouraud broth (Difco, Detroit, Michigan) and its exo-antigens concentrated by ultrafiltration under positive pressure in a stirred cell fitted with a PM-10 membrane (Amicon Corp., Lexington, Massachusetts).

Agar gel double diffusion was carried out in plastic petri dishes (100 by 15 mm) into which 7.5 ml of 0.25% phenolized-1% purified agar (Difco) was added. A seven-well pattern with 4-mm-diameter wells 4 mm apart was used. The antigen was added into the central well and positive control serum was placed into the top and lower wells. The testing sera were added into the four remaining wells.

The ELISA was carried out as per Mendoza et al. (Mendoza et al., Clin. Diag. Lab. Immunol. 4: 715-718 (1997)). Flat-bottom polystyrene microtiter plates (96-well Immulon2; Dynatech Laboratories Inc., McLean, Virginia) were coated with the antigen prepared as above and incubated overnight at 4° C and then blocked for 1 h at 37° C with 5% gelatin. Dilutions of the sera

under investigation were prepared and then added to the coated plates and incubated for 1 h. After several washes, 100 µl of a horseradish peroxidase-conjugated rabbit anti-horse, anti-dog, or anti-cat (heavy and light chains) antibody was added to each well and then incubated at 30° C for 1 h. After incubation, the reaction was stopped with chromogen buffer and color development was recorded in a Dynatech MR 5000 ELISA plate reader at 490 nm. The immunoperoxidase assay performed on sera from two of the dogs was not carried out in our facilities, but was carried out in other laboratories at the request of their owners.

Due to the presence of large quantities of eosinophils in the infected tissues of horses with pythiosis, an anti-horse-IgE ELISA was also developed to investigate the titers of this immunoglobulin before and after vaccination with the PIV. The anti-horse-IgE peroxidase-conjugated immunoglobulin was produced by Bio-Medical International (Austin, Texas). In brief, 50 µl of the diluted horse serum sample were added to each well and then incubated at room temperature for 2 hrs. After several washes, the plates were blocked with 5% gelatin. The plates were then washed and 50 µl of a 1/10,000 dilution of the anti-horse-IgE peroxidase-conjugated was added to each well and then incubated at room temperature for two hrs. Color development and titer readings were according to the ELISA previously described. An ELISA-IgE using antigen of *Pythium insidiosum* was only tested in horses.

Vaccine Production and Administration. For the experimental use of the PIV in animals with pythiosis, a FDA permit was requested and granted to Bio-Medical International. The PIV was prepared as follows. In brief, *Pythium insidiosum* (ATCC 74446) was grown in Sabouraud broth at 37° C. The liquid growth medium containing the extracellular antigens was removed from the cell mass and saved. The cell mass was washed twice

with sterile distilled water and then disrupted by sonication. The resulting supernatant containing the intracellular antigens was separated from the pellet containing the disrupted cell debris and mixed with the liquid growth medium containing the extracellular antigens. The extracellular antigens and the intracellular antigens in the mixture were precipitated with acetone and the precipitate resuspended in water to produce the PIV with a protein concentration of about 2.0 µg/ml. The PIV was then lyophilized and stored at -80° C until used. The same PIV batch was used throughout the study.

The PIV was administered at least twice in all animals. The first vaccination was given on day 0 and the second given on day fifteen. Briefly, 100 µl of the PIV was intradermally injected in the middle of the neck of the animals on day 0. The second injection was subcutaneously administered fifteen days later. Additional injections were given subcutaneously to animals that did not show improvement with the first two vaccinations. Based in previous studies (Alfaro et al., Equine Vet. J. 22: 295-297 (1990); Miller, Aust. Vet. J. 67: 377-82 (1981); Mendoza et al., Mycopathologia 119: 89-95 (1992)), the inflammatory reaction at the site of injection was recorded as weak for indurations less than 29 mm in diameter, mild for indurations about 30 to 99mm in diameter and strong for indurations greater than 100 mm in diameter. The animals were recorded as cured if all traces of the pythiosis-induced skin lesions had been eliminated. In animals with skin lesions this included: epithelialization of the lesion, cessation of discharge, and closure of the sinus tracts. The dog with intestinal involvement was evaluated by monitoring the decline of its clinical symptoms such as vomiting and diarrhea as well as the reduction in size of the tumor-like intestinal masses as determined by X-ray and palpation.

Diagnosis of Pythiosis and Histopathological Studies. Pythiosis-induced lesions in the horses were more commonly observed around their limbs, abdomen, inguinal areas, face, and shoulders, respectively (Table 1). The four dogs were diagnosed with subcutaneous pythiosis and one had intestinal pythiosis. The cat presented a non-ulcerative subcutaneous lesion on the chest. All the animals had been unsuccessfully treated either by surgery or chemotherapy (Ivermectin, steroids, antibiotics, or antifungals, including itraconazole, and others) prior to immunotherapy. In all cases, the diagnosis of pythiosis in the animals was confirmed by serological methods, and for several of the animals, by serology plus histopathology and immunoperoxidase assay. The serological assays established that all animals had detectable anti-*Pythium insidiosum* IgG titers of 1:3200 or greater. The ID showed that all animals with clinical lesions had identity bands with the positive control sera. All of these parameters were consistent with pythiosis. In the IgE-ELISA, titers of 1:6,400 and greater were detected in animals with active pythiosis. The IgG and IgE titers declined in cured cases and were almost undetectable six months after successful immunotherapy.

Histopathological studies of the infected tissues showed an inflammatory response with numerous eosinophils, mast cells, neutrophils, plasma cells, and macrophages in the infected species (Figure 1). Silver-stained samples from the horses with pythiosis showed that the hyphae of *Pythium insidiosum* were restricted to small calcareous-like masses termed "kunkers," a typical feature of equine pythiosis. Hyphal elements of *Pythium insidiosum* could not be detected outside of these kunkers. The dog and cat evaluated by histopathology did not show these kunkers. In these animals, *P. insidiosum*'s hyphae were found in the center of the heavily eosinophilic reactions.

Table 1. Clinical features of equine cases with pythiosis used in this study and their responses to *Pythium insidiosum*-vaccine

State	Age/sex	Lesions	Duration of illness	Diagnosis	Previous treatments	Vaccination reaction	Outcome
AR (Gi)	4 y/F	Abdomen, 220X220 mm	4 months	ID, ELISA (+)	Surgery, drugs	Strong, 123 mm	Cured
FL (Sc)	13 y/F	Face, 80X30 mm	>4 months	ID, ELISA (+)	Surgery, Drugs	Mild, 25 mm	Cured
FL (Sn)	12 y/M	Limb, 150X100 mm	> 2 months	ID (+), clinical	Surgery	Mild, 60 mm	Cured
FL (Jo)	20 y/M	Limb, 300X150 mm	> 2 months	ID (+), clinical	Several surgeries	Mild, 60 mm	Cured
FL (Wa)	8 y/F	Limb, 60X50 mm	5-7 days	ID (+), Clinical	Surgical debridement	Strong, 100 mm	Cured
FL (Ho)	5 y/F	Shoulder & abdomen 50X50 & 120X20mm	15 days	ELISA (+)	Cryosurgery	Strong, 200 mm	Cured
FL (Pe)	22 y/M	Limb, 60X40	17 days	ID, ELISA (+)	Surgical debridement	Mild, 30 mm	Cured
LA (Re)	5 y/M	Limb, 120X100 mm (two lesions)	2 months	ELISA (+)	Surgical debridement	Mild, 90 mm	Cured*
MS (Ba)	3 y/F	Limb, 250X250 mm	>2 months	ELISA (+)	Topical drugs	Strong, 150 mm	Cured***
MS (Pa)	15 y/F	Limb, 300X200 mm	4 months	ELISA (+)	Topical drugs	Strong, 170 mm	Cured
MS (Im)	2 y/M	Limb, 300X300 mm	4 months	ELISA (+)	Topical drugs	Strong, 200 mm	(Vaccine + surgery) Cured
MS (So)	7 y/F	Limb, 240X240 mm	3 months	ELISA (+), Histopathology (+)	Topical Drugs	Strong, 150 mm	(Vaccine + surgery) sacrificed (†)
NC (Sa)	20 y/M	Inguinal, 60X50 mm (two lesions)	>2 months	ID (+), ELISA (+)	Surgery	Mild, 40 mm	Not cured
NC (Ga)	14 y/F	Inguinal 200X150 mm	>2 months	ID (+), ELISA (+)	Surgery	Mild, 30 mm	Not cured
TN (Re)	4 y/F	Abdomen, 200X80 mm	2 months	ELISA (+)	Topical Drugs	No response	Not cured
TX (Ta)	13 y/M	Limb, 250X100 mm	~4 months	ELISA (+)	Topical Drugs	Strong, 200 mm	Died††
TX (Ah)	5 y/M	Limb, 280X210 mm	1 month	Clinical, kunkers ELISA (+)	Surgery, drugs	Mild, 30 mm	Cured
TX (Co)	6 y/F	Limb, 100X80 mm	1 month	ELISA (+)	Topical drugs§§	Weak, 15 mm	Not cured
TX (Sn)	22 y/M	Mouth, 150X100 mm	>2 months	ELISA (+) Histopathology (+)	Surgical debridement	Weak, 5 mm	Not cured

Serial biopsies in a horse cured after immunotherapy showed that the initial eosinophilic response was replaced by mononuclear macrophages and cytotoxic T lymphocytes (CTL) (Figure 1). Silver-stained samples in the serial biopsies revealed that the hyphae slowly became damaged by the cell's mediated mononuclear immune response (Figure 1). In the early stages of cure, seven days into successful immunotherapy, numerous ghost hyphal cells were readily observed in the healing tissues. Hyphal elements of *Pythium insidiosum* could not be detected 15 days after the first injection in successfully cured horses (Figure 1).

Responses of the Infected Animals to Immunotherapy. Injected animals showed different sizes of indurations at their injection sites. Of the 19 horses tested, one did not react to the injection and was not cured, two showed weak indurations at the injection site of 15 mm or less in diameter but did not respond to immunotherapy, eight developed mild reactions with indurations at the injection site between 30 to 90 mm in diameter, and eight had strong inflammatory reactions with indurations at the injection site greater than 100 mm in diameter (Table 1). Seven of the eight horses with a mild inflammatory reaction were successfully cured. The remaining horse with a mild inflammatory reaction showed an initially curative response but the infection recurred and could not be cured even after several injections of the PIV. Anaphylactic reactions to the PIV were not detected in the animal, even after more than 12 injections of the PIV. All horses with a strong inflammatory reaction at the injection site were cured. However, two of the horse died during this study of causes not related to this trial or pythiosis (Table 1).

Two of the five dogs with the disease strongly reacted to the PIV at the injection site, but only one dog was cured. One of the three remaining dogs had a

weak inflammatory reaction at the injection site and the other two dogs did not have any inflammatory reaction. None of the dogs were cured of pythiosis. The cat developed a weak inflammatory reaction at the injection site but was not cured of pythiosis (Table 2).

The Mendoza vaccine tested in Australia and Costa Rica cured about 57% of all injected horses with pythiosis (Miller, Aust. Vet. J. 67: 377-82 (1981); Mendoza et al., Mycopathologia 119: 89-95 (1992)); however, the Mendoza vaccine failed to cure chronically infected horses. The failure of the Mendoza vaccine to cure chronically infected horses of pythiosis was related to the immunological status of the vaccinated horses not because the Mendoza vaccine was inadequate (Mendoza et al., Mycopathologia 119: 89-95 (1992)). In (Mendoza et al., Mycopathologia 119: 89-95 (1992)), 100% of the equines of less than 20 days infection with pythiosis were cured after immunotherapy. However, the efficacy of the Mendoza vaccine diminished in horses which had pythiosis for one month and the Mendoza vaccine failed in 100% of the horses chronically infected with pythiosis (pythiosis infection greater than 2 months duration). A modification of the Mendoza vaccine (the PIV of the present invention) was introduced in 1998 (Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)). This modified Mendoza vaccine (PIV) cured four of seven horse pythiosis infections of more than four months duration (70% effectiveness), as well as all of the horses with acute pythiosis. The data presented in this example with PIV is in agreement with their preliminary analysis.

Although Mendoza et al. (Mendoza et al., Mycopathologia 119: 89-95 (1992)) suggested that the efficacy of the PIV depends mainly on the immunological status of the infected hosts, the addition of *Pythium insidiosum*'s cytoplasmic antigens to the Mendoza vaccine

Table 2. Clinical features of dogs and a cat with pythiosis used in this study and their responses to *Pythium insidiosum*-vaccine

Species	Age/sex	Lesions	Duration of illness	Diagnosis	Previous treatments	vaccination reaction	Outcome
Dog, FL (Ho)	4 y/F	Limbs (two lesions) ~100X50 mm	>2 months	ID (+), ELISA (+)	Surgery	Strong, 250 mm	cured
Dog, FL (Pr)	4 y/F	Perianal 30X20 mm	>5 months	Histopathology (+) ID (+)	Surgery	No response	Not cured
Cat, FL (Ra)*	3 y/M	Non-ulcerative granuloma (chest) 120X50 mm	1 year	Histopathology (+) ID (+)	Drugs	Weak, 5 mm	Not cured
Dog, LA (MA)	3y/M	Intestinal granuloma	>3 months	Immunoperoxidase (+)	Antibiotics	Strong, 250 mm	Not cured
Dog, MS (Ma)	1 y/F	Rump and Tail 80X70 mm	5 months	Immunoperoxidase (+)	Antifungals	No response	Not cured
Dog, NC ()	2y/F	Limbs 35X80 mm	4 months	ID (+)	Antifungals	Weak, 7 mm	Not cured

* This case was also cited in reference 23

as described herein (Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)) indicates that, in addition to a sound immune system, the PIV's immunotherapeutic properties is also affected by the type of immunogens used to manufacture the vaccine. For example, Mendoza et al. (Mendoza et al., Mycopathologia 119: 89-95 (1992)) found that horses with chronic pythiosis were not cured by a vaccine made of only *Pythium insidiosum*'s extracellular antigens (Mendoza vaccine).

In contrast, the data presented herein and in previous studies (Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)), show that the PIV of the present invention, containing an equal mixture of extracellular antigens and intracellular antigens (cytoplasmic immunogens), was able to cure about 73% of the 26 injected horses including chronically infected horses. This indicates that the addition of the intracellular antigens to the Mendoza vaccine dramatically increased the efficacy of the Mendoza vaccine. As in previous trials using the Mendoza vaccine, the anergic cases did not react to the PIV as well. Thus, it appears that the immunotherapeutic properties of pythiosis vaccines are affected by the immunogens used to manufacture the PIV as well as by the immunological status of the infected host.

Interestingly, only one of the five canines (20%) with pythiosis responded to the PIV and the only cat used in this study did not react at all. It had been found that horses with a weak or no reaction at the injection site were not cured (Alfaro et al., Equine Vet. J. 22: 295-297 (1990); Mendoza et al., Mycopathologia 119: 89-95 (1992)). The horses in those groups were always found to be anergic. The finding could explain the failure of the PIV in four of the six animals in Table 2, but it does not explain why one of

the two dogs with strong inflammatory reactions did not respond to the immunotherapy. The findings in dogs are in contrast with the data in horses. For instance, horses with strong reactions at the site of injection all responded to immunotherapy (Newton et al., Compendium. 15: 491-493 (1993); Mendoza et al., Mycopathologia 119: 89-95 (1992); Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998)).

The poor performance of the PIV in dogs and cats contrasts with the high number of cured horses and its excellent performance in humans with arterial pythiosis (Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)). At least four of the five humans with pythiosis (80% effectiveness) were cured with the PIV (unpublished data). The failure of the PIV to cure one of the two dogs with strong inflammatory reactions at the injection site is intriguing and merits additional studies.

Serial biopsies, taken in successfully injected horses, showed that the eosinophils were gradually replaced by CTL-like cells and mononuclear macrophages (Figure 1). Three to six days after injection, a mononuclear reaction was evident around the hyphal elements of *Pythium insidiosum*. Degenerating hyphae of *Pythium insidiosum* were observed by day seven and hyphal elements of *Pythium insidiosum* could not be detected 15 days after the first PIV injection. Interestingly, healing of the cutaneous lesions was always noticeable seven days after PIV injection which coincided with the disappearance of hyphae from the tissues. Our data and that of others (Julia et al., Science 274: 421-3 (1996); Haberer et al., Infect. Immunol. 66: 3100-05 (1998); Secrist et al., J. Exp. Med. 178: 2123-30 (1993)) suggest that the modulation of cell mediated responses and the release of other immune-active compounds are the driving force behind the PIV's immunotherapeutic properties.

The changes in cell mediated responses before and after PIV injection in the horses of this study and those in humans cured by the PIV (Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)), further support the concept of immune-modulation in the cured horses. Recent studies have found that changes in cytokine profiles can modulate a host's immune response to antigens (Steidler et al., Science 289: 1352-55 (2000); Convit et al., Trans. R. Soc. Trop. Med. Hyg. 87: 444 (1993)). For instance, human patients with pythiosis always display interleukin 4 (IL4) (Mendoza et al., J. Clin. Microbiol. 25: 2159-2162 (1987)). In contrast, interleukin 2 (IL2) is more prominent in humans successfully treated by immunotherapy. In addition, high titers of IgE are present in humans with pythiosis. High titers of IgE are also found in horses with pythiosis. However, the IgE titers in the horses declined after three months of successful PIV immunotherapy, a finding that parallels that found in humans successfully treated by immunotherapy (Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Mendoza et al., J. Clin. Microbiol. 25: 2159-2162 (1987)). Since eosinophils and IgE are commonly associated with a Th2 response, their presence before immunotherapy and subsequent disappearance after successful vaccination strongly supports the concept of switching in the T helper subsets.

Earlier studies have shown that the Th1 subset, which produces IL2 and interferon γ (INF γ), is involved in the clonal expansion of CTL, mononuclear macrophage activation, and production IgG isotypes which mediate complement activation against sensitized pathogens (15; Steidler et al., Science 289: 1352-55 (2000); Convit et al., Trans. R. Soc. Trop. Med. Hyg. 87: 444 (1993)). The Th2 subset produces IL4 and interleukin 5 (IL5), which stimulate neutralizing antibodies and IgE, the initiators of hypersensitivity

and eosinophilia (Cock et al., J. Clin. Microbiol. 25: 344-349 (1987); Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998)). Based on the cellular events before and after successful immunotherapy in humans and horses with pythiosis, one can theorize that switching from a Th2 immune response before vaccination (eosinophils, mast cells, and the presence of IgE, IL5), to Th1 immunity (mononuclear macrophages, CTL, IL2), may be the driving factor in the curative properties of the PIV.

This is further supported by the finding that *Pythium insidiosum* possesses two different sets of antigens that seem to stimulate different subsets of T helper cells. One of these antigens are extracellular antigens expressed *in vivo* and in cultures (Mendoza et al., J. Clin. Microbiol. 31: 2967-73 (1993)). The extracellular antigens stimulate a topical allergic reaction with eosinophils and IgE when injected into horses and rabbits (unpublished data). In contrast, the cytoplasmic antigens have been credited with curative properties (Miller, Aust. Vet. J. 67: 377-82 (1981); Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998), Mendoza et al., J. Clin. Microbiol. 25: 2159-2162 (1987)). The detection of secreted extracellular antigens within kunkers during natural horse infection using fluorescent antibodies strongly supports this concept (Mendoza et al., J. Clin. Microbiol. 31: 2967-73 (1993)). Thus, *Pythium insidiosum* expresses a soluble extracellular antigen or antigens during natural infections that triggers eosinophils, mast cells, IgE, and possibly IL4 and IL5. The extracellular antigen(s) appear to lock the immune response in a Th2 mode, which in turn is responsible for clinical features of the disease. Because the cytoplasmic antigens seem to stimulate the host's immune system into Th1 curative immunity, the enhancement of the efficacy of Mendoza's vaccine by the addition of cytoplasmic antigens to the Mendoza vaccine in this and another study (Dixon et al.,

Med. Mycol. 36(Supp. 1): 57-67 (1998)) strongly supports this idea.

Based on these data, a hypothesis that explains the pathogenesis of pythiosis as well as the immunotherapeutic mechanisms of the PIV is proposed as shown in Figure 2. As shown in Figure 2, pythiosis begins when a host comes in contact through an open wound with propagules of *Pythium insidiosum* (usually a zoospore as shown in the diagram, but it could be any hyphal element). A germ tube is then formed from the propagules which mechanically penetrates the tissue of the host (brown line). Once in the tissue, *Pythium insidiosum*'s hyphae release extracellular antigens (brown dots) that are exposed to antigen-presenting cells (APC). The APC release IL4 that drive naive T helper cells (Th0) into the Th2 subset which in turn produces more IL4 and IL5. IL4 down regulates the Th1 subset (dotted light brown arrows) and stimulates B cells to produce IgE, IgM, and precipitin IgG used for diagnostic purposes. The IL5 and IgE trigger production of eosinophils and mast cells at the site of injury (purple arrows). These cells then degranulate (responsible for the tissue damage) over *Pythium insidiosum*'s hyphae triggering the Hoeppli-Splendore phenomenon, which later evolves into kunkers (only in equine pythiosis) (blue arrow). *Pythium insidiosum* multiplies inside the kunkers where it will produce overwhelming quantities of extracellular antigens, an event that ultimately locks the immune response into a Th2 mode.

Also shown in Figure 2 is when PIV's cytoplasmic antigens are injected into the host with pythiosis (upper right side), the APC are presented, in a different fashion, the cytoplasmic antigens which in the natural infection, the immune system did not properly recognize. The APC releases interferon gamma (INF γ) that drives naive Th0 into the Th1 subset. The

Th1 subset produces more IL2 and INF γ , in small quantities at first, because the Th1 subset had been down regulated by the IL4 released from the Th2 subset. In turn, IL2 and INF γ trigger a mononuclear cell mediated immune (CMI) response composed mostly by cytotoxic T lymphocytes (CTL) and macrophages which damage and destroy the hyphae of *Pythium insidiosum* (red arrows). It is not clear if during immunotherapy IL2 and INF γ also stimulate B cells to produce protective IgG classes. The production of INF γ at the site of infection by the APC and the Th1 subset results in the down regulation of the Th2 subset (dotted pink arrow). The down regulation of Th2 and the fact that the hyphae, which have been damaged by the CMI, no longer produce extracellular antigens to lock the host's immune system in the Th2 mode may explain why horses and humans with pythiosis are cured by the PIV.

Therefore, in view of the hypothesis, the propagules of *Pythium insidiosum* contact the host (The Jordan Report. Accelerated Development of Vaccines. Division of Microbiology and Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institute of Health (NIH), Bethesda, MD. p. 17 (2000)). Upon contact through an open wound, *Pythium insidiosum*'s propagules form a germ tube that mechanically penetrates tissues. Once into the tissue, *Pythium insidiosum*'s hyphae produce extracellular antigens (Mendoza et al., J. Clin. Microbiol. 31: 2967-73 (1993)) which trigger a Th2 immune response and the production of eosinophils, mast cells, IgE, IL4, and IL5. The constant production of the extracellular antigens causes the host's immune response to become locked into a Th2 mode. The overwhelming number of degranulated eosinophils (Hoepli-Splendore phenomenon) and mast cells around the hyphae of *Pythium insidiosum* are primarily responsible for the extensive and rapid tissue damage encountered during pythiosis, especially

in horses and dogs (Mendoza et al., J. Mycol. Med. 6: 151-164 (1996); Chaffin et al., Vet. Clin. North. Am. Equine Pract. 11: 91-103 (1995)).

Our data and data of others (Newton et al.,
5 Compendium. 15: 491-493 (1993); Miller, Aust. Vet. J. 67: 377-82 (1981); Mendoza et al., Mycopathologia 119: 89-95 (1992); Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Thitithanyanont et al., Clin. Infec. Dis. 27: 1394-400 (1998)), suggest that the production of the
10 Hoeppli-Splendore reaction and the secretion of the extracellular antigens are evolutionary strategies developed by *Pythium insidiosum* to secure its proliferation in the host's tissues. This is supported by the observation that viable hyphae of *Pythium*
15 *insidiosum* have been found only inside the eosinophilic reaction (Mendoza et al., J. Mycol. Med. 6: 151-164 (1996)), or inside the kunkers in equine pythiosis (Alfaro et al., Equine Vet. J. 22: 295-297 (1990); Chaffin et al., Vet. Clin. North. Am. Equine
20 Pract. 11: 91-103 (1995); Mendoza et al., Mycopathologia 119: 89-95 (1992)), indicating that *Pythium insidiosum* might use the Hoeppli-Splendore phenomenon for its survival.

Conversely, it has been shown that injecting
25 the cytoplasmic immunogens of *Pythium insidiosum* stimulates a mononuclear response in the host that cures the host or in some instances, protects the host for a short period of time (Dixon et al., Med. Mycol. 36(Supp. 1): 57-67 (1998); Thitithanyanont et al., Clin. Infec.
30 Dis. 27: 1394-400 (1998); Mendoza et al., J. Clin. Microbiol. 25: 2159-2162 (1987)). The findings strongly suggest that upon contact with the cytoplasmic antigens, a down-regulation of the Th2 response is triggered. In our hypothesis (Figure 2), the injected cytoplasmic
35 antigens are displayed to the host's immune system in a different fashion than during natural infection. It is the different presentation of the cytoplasmic antigens

which stimulates the Th1 response. The observation that in successfully vaccinated humans and animals the IgE and IL4 levels decline, the IL2 and INF γ levels increase, and the eosinophils are replaced by CTLs and macrophages (Mendoza et al., J. Clin. Microbiol. 25: 2159-2162 (1987)), supports the hypothesis.

Because it has been shown that INF γ down regulates the Th2 subset and that the Th1 response triggers a cell mediated immune response that eliminates pathogens from infected tissues, we further hypothesize that the hyphae of *Pythium insidiosum* are killed by the mononuclear response triggered by the up-regulated Th1 subset. The up-regulated Th1 induces the production of CTLs and macrophages which kill the hyphae thereby preventing production of the extracellular antigens required to maintain the Th2 response.

Therefore, the activation of a Th1 subset by the cytoplasmic antigens and the subsequent down regulation of the Th2 response, could well explain the PIV's curative properties. Similar mechanisms have been reported for immunotherapeutic treatments against other eukaryotic pathogens (Julia et al., Science 274: 421-3 (1996); Haberer et al., Infect. Immunol. 66: 3100-05 (1998); Convit et al., Trans. R. Soc. Trop. Med. Hyg. 87: 444 (1993)) and to control allergic reactions (Secrist et al., J. Exp. Med. 178: 2123-30 (1993); Steidler et al., Science 289: 1352-55 (2000)).

Until recently, pythiosis was considered an exotic disease of equines and canines in the United States (Thomas et al., Compendium 20: 63-75 (1998)). A recent increase in the number of reported cases of pythiosis suggests that the disease is more prevalent than previously suspected. This upsurge in the number of reported cases may be related to the development of new diagnostic tools (Mendoza et al., J. Clin. Microbiol. 23: 813-816 (1986); Mendoza et al., Clin. Diag. Lab. Immunol. 4: 715-718 (1997); Brown et al., Am.

Vet. Med. Res. 49: 1866-68 (1988)) and by the increased awareness of clinicians to *Pythium insidiosum*'s presence in enzootic areas. The majority of these reports have come from well known enzootic areas of pythiosis in such states as Alabama, Arkansas, Florida, Georgia, Illinois, Louisiana, Mississippi, Missouri, North and South Carolina, Oklahoma, Tennessee, and Texas (Mendoza et al., J. Mycol. Med. 6: 151-164 (1996); Chaffin et al., Vet. Clin. North. Am. Equine Pract. 11: 91-103 (1995)). However, reports are also coming in from previously unsuspected areas including California, New Jersey, Pennsylvania, Wisconsin and New York. The numerous canine and equine cases found during our studies and the increasing reports of dog pythiosis (Mendoza et al., J. Mycol. Med. 6: 151-164 (1996); Thomas et al., Compendium 20: 63-75 (1998)) strongly suggest that this disease can no longer be considered a rare clinical entity of equines and canines in North America.

EXAMPLE 7

This example illustrates the method for evaluating the therapeutic and prophylactic properties of the PIV in a rabbit model. The rabbit model provides information on why the PIV failed in dogs or other animals with chronic pythiosis, how the PIV can be improved, and unveil the PIV's prophylactic and therapeutic attributes.

Pythium insidiosum type strain (ATCC 74446) is used to challenge rabbits and prepare PIV as in Example 1. Briefly, Strain ATCC 74446 is transferred to a one liter flask containing 500 ml of Sabouraud dextrose broth. The inoculated flask is incubated at 37° C for five days on a rotary shaker at 150 rpm. The cultures are then killed with Merthiolet (0.02% w/v) and filtered to separate the cells (hyphae) from the liquid phase containing the secreted extracellular antigens. The liquid phase is stored in a sterile container. The mass

of cells retained by the filter is washed twice with distilled water, disrupted by sonication until 100% of the hyphae are fragmented, and then centrifuged at 5,000 x g for 20 minutes. The supernatant fraction containing the intracellular cytoplasmic antigens is separated from the cell debris pellet and added to the stored liquid phase. The intracellular and extracellular antigens in the mixture is precipitated with an equal volume of acetone and pelleted by centrifugation at 20,000 x g for 30 minutes. The pellet is resuspended in sterile distilled water to a final protein concentration of about 2 mg/ml and dialyzed in sterile distilled water using a membrane with a 10,000 MW cut-off to produce the PIV. Sterility of the PIV is confirmed by culturing aliquots of the PIV on blood agar and Sabouraud dextrose broth. PIV is stored at 4° C until use. Placebo vaccines are made as above except that the ATCC 74446 strain is not added to the culture flask.

Rabbits are infected with *Pythiosum insidiosum* zoospores. Zoospores are used as the infecting propagules because they are easy to prepare and can be easily counted. This ensures consistent challenge doses. Furthermore, 100% of the rabbits injected with zoospores developed subcutaneous nodules or systemic infections depending on the route of injection (Mendoza and Prendas, Mycopathologia 104: 59-62 (1988)). Thus, zoospores are effective propagules to reproduce cutaneous and systemic experimental pythiosis in rabbits. To induce zoospore production in the laboratory, the methodology of Mendoza and Prendas, Mycopathologia 104: 59-62 (1988), is used. The zoospores are counted and standardized to about 50 zoospores/100 µl (about 500 zoospores/ml). This is the minimum quantity of zoospores required to produce cutaneous and systemic infections in rabbits (Mendoza and Prendas, Mycopathologia 104: 59-62 (1988)).

To evaluate the prophylactic response of the

PIV in the rabbit model, rabbits are vaccinated with PIV on days 0 and 15 and followed by subcutaneous injection with viable zoospores on days 20 and 60 after the first vaccination. It was previously found that rabbits produced anti-*Pythium insidiosum* IgG around 25 to 30 days post-vaccination (Reyes et al., An animal model to study *Pythium insidiosum* infections. 14th Meeting of the International Society for Human and Animal Mycology, Buenos Aires, Argentina, p. 225 (2000)), which indicated that the immune response was activated at about that time. Control rabbits are injected with the placebo. Protection is measured by the lack of cutaneous granulomatous lesions at the site of challenge.

To evaluate the therapeutic response of the PIV in infected rabbits, rabbits which have been first infected with the zoospores and have developed subcutaneous pythiosis, are vaccinated with the PIV. Rabbits are recorded as cured if all traces of the lesions have been eliminated. This includes epithelization of the lesion, cessation of discharge, and closure of the sinus tracts.

Evaluation of the rabbit's immune response after experimental infection and after PIV immunization is as follows. Immunodiffusion (ID), ELISA, and Western blots are used to evaluate the rabbit's antibody levels after PIV vaccination and during experimental pythiosis. The ID test is used to show sero-conversion in immunized and challenged rabbits. ELISA is used to evaluate the titers of IgG in vaccinated and challenged rabbits. Western blots is used to determine the presence of antibodies against the immunogenic antigens of *Pythium insidiosum*. In addition, IgG isotype assays, cytokine assays, and leukocyte counts are used to evaluate the rabbit's immune response to the PIV in all of the vaccinated and challenged rabbits. Leukocyte counts are performed on the rabbit's total blood using a cell counter chamber. All rabbits in the prophylactic

experiments are bled before immunization or before zoospore challenge and sera stored at -80° C until tested. Similarly, rabbits in the therapeutic experiments are bled before zoospore challenge to evaluate the humoral immune response before infection and 20 days after PIV injection.

To evaluate the different IgG isotypes triggered in the experimentally-induced pythiosis or by the PIV, and IgG isotype assay is performed. Rabbits are bled before inducing pythiosis, 14 days post-vaccination, and 14 days after the second immunization. The isotype assays measure the total immunoglobulin populations in the rabbit. briefly, 50 µl of the PIV (2 mg/ml) is coated on flat-bottomed polystyrene microtiter plates (96-well, Immulon 2, Dynatech Laboratories, Inc., Virginia) at 4° C for 24 hr. The plates are then reacted against the rabbit sera as per Mendoza et al., Clin. Diagn. Lab. Immunol. 4: 715-718 (1997) followed by reacting with anti-IgG isotypes (IgG₁, IgG₂, and IgG₃) according to the manufacturer's instructions (Accurate Chemicals, New York). In addition, to monitor the Th2 to Th1 switching, IgE levels in all the rabbits are determined. The immunological data provided by the method provides a means for validating the Th2 to Th1 switching hypothesis which can be extrapolated to other infectious diseases of animals.

The method for evaluating the prophylactic and therapeutic properties of the PIV is of importance for preventing or curing pythiosis for all animals inhabiting enzootic areas of pythiosis. Until now, the assessment of the PIV's therapeutic and prophylactic properties was done using clinical cases of the disease in equines and in humans. However, cases of pythiosis in horses are usually far away and the experimental conditions outside the laboratory are in general, not controllable. Therefore, the method for evaluating the prophylactic and therapeutic properties of the PIV in

the rabbit model system is of profound importance for preventing or treating pythiosis in animals.

While the present invention is described herein with reference to illustrated embodiments, it
5 should be understood that the invention is not limited hereto. Those having ordinary skill in the art and access to the teachings herein will recognize additional modifications and embodiments within the scope thereof. Therefore, the present invention is limited only by the
10 claims attached herein.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218